

Physiological —— and —— Molecular Plant Pathology

EDITORS

J. P. Blakeman, W. R. Bushnell, D. D. Clarke, M. J. Daniels, R. Hammerschmidt,
D. R. Walters

EDITORIAL BOARD

P. Albersheim	R. S. S. Fraser	G. Loebenstein	G. F. Pegg
J. A. Bailey	B. J. M. Fritig	K. Mendgen	L. C. van Loon
T. Boller	A. Fuchs	D. Mills	E. W. B. Ward
L. D. Dunkle	M. C. Heath	S. Ouchi	O. C. Yoder
A. H. Ellingboe	C. J. Lamb		

Volume 41, 1992

ACADEMIC PRESS

Harcourt Brace & Company, Publishers

London San Diego New York Boston Sydney Tokyo Toronto

Copyright © 1992 Academic Press Ltd.

ALL RIGHTS RESERVED

No part of this volume may be reproduced in any form,
by photostat, microfilm, or any other means, without
written permission from the publishers.

Contents of Volume 41

Number 1, July 1992

PIERPOINT, W. S., GORDON-WEEKS, R. and JACKSON, P. J., The occurrence of the thaumatin-like, pathogenesis-related protein, PR-5, in intercellular fluids from <i>Nicotiana</i> species and from an interspecific <i>Nicotiana</i> hybrid	1
GOY, P. A., FELIX, G., MÉTRAUX, J. P. and MEINS, F. JR., Resistance to diseases in the hybrid <i>Nicotiana glutinosa</i> × <i>Nicotiana debneyi</i> is associated with high constitutive levels of β -1,3-glucanase, chitinase, peroxidase and polyphenoloxidase	11
WUBBEN, J. P., JOOSTEN, M. H. A. J., VAN KAN, J. A. L. and DE WIT, P. J. G. M., Subcellular localization of plant chitinases and 1,3- β -glucanases in <i>Cladosporium fulvum</i> (syn. <i>Fulvia fulva</i>)-infected tomato leaves.. .. .	23
BENHAMOU, N. and THERIAULT, G., Treatment with chitosan enhances resistance of tomato plants to the crown and root rot pathogen <i>Fusarium oxysporum</i> f. sp. <i>radicis-lycopersici</i>	33
PASCHOLATI, S. F., YOSHIOKA, H., KUNOH, H. and NICHOLSON, R. L., Preparation of the infection court by <i>Erysiphe graminis</i> f. sp. <i>hordei</i> : cutinase is a component of the conidial exudate	53
RICKER, K. E. and BOSTOCK, R. M., Evidence for release of the elicitor arachidonic acid and its metabolites from sporangia of <i>Phytophthora infestans</i> during infection of potato	61
ERRATUM.. .. .	73

Number 2, August 1992

RANSOM, RICHARD F., HIPSKIND, JOHN, LEITE, BRENO, NICHOLSON, RALPH L. and DUNKLE, LARRY D., Effects of elicitor from <i>Colletotrichum graminicola</i> on the response of sorghum to <i>Periconia circinata</i> and its pathotoxin	75
ESSENBERG, MARGARET, PIERCE, MARGARET L., HAMILTON, BETTY, COVER, ELLEN C., SCHOLES, VERNON E. and RICHARDSON, PAUL E., Development of fluorescent, hypersensitively necrotic cells containing phytoalexins adjacent to colonies of <i>Xanthomonas campestris</i> pv. <i>malvacearum</i> in cotton leaves	85
ESSENBERG, MARGARET, PIERCE, MARGARET L., COVER, ELLEN C., HAMILTON, BETTY, RICHARDSON, PAUL E. and SCHOLES, VERNON E., A method for determining phytoalexin concentrations in fluorescent hypersensitively necrotic cells in cotton leaves	101
PAIN, N. A., O'CONNELL, R. J., BAILEY, J. A. and GREEN, J. R., Monoclonal antibodies which show restricted binding to four <i>Colletotrichum</i> species: <i>C. lindemuthianum</i> , <i>C. malvarum</i> , <i>C. orbiculare</i> and <i>C. trifolii</i>	111
CASTORIA, R., FANELLI, C., FABBRI, A. A. and PASSI, S., Metabolism of arachidonic acid involved in its eliciting activity in potato tuber	127
RGHEI, N. A., CASTLE, A. J. and MANOCHA, M. S., Involvement of fimbriae in fungal host-mycoparasite interaction	139

Number 3, September 1992

CARVER, T. L. W., ROBBINS, M. P., ZEYEN, R. J. and DEARNE, G. A. Effects of PAL-specific inhibition on suppression of activated defence and quantitative susceptibility of oats to <i>Erysiphe graminis</i>	149
LEE-STADELMANN, O. Y., CURRAN, C. M. and BUSHNELL, W. R. Incompatibility conditioned by the <i>Mla</i> gene in powdery mildew of barley: change in permeability to non-electrolytes	165
XUE, B., GOODWIN, P. H. and ANNIS, S. L. Pathotype identification of <i>Leptosphaeria maculans</i> with PCR and oligonucleotide primers from ribosomal internal transcribed spacer sequences	179
CORDERO, M. J., RAVENTÓS, D. and SAN SEGUNDO, B. Induction of PR proteins in germinating maize seeds infected with the fungus <i>Fusarium moniliforme</i>	189
FLOOD, J., WHITEHEAD, D. S. and COOPER, R. M. Vegetative compatibility and DNA polymorphisms in <i>Fusarium oxysporum</i> f.sp. <i>elaeidis</i> and their relationship to isolate virulence and origin	201
ALLAN, R. H., THORPE, C. J. and DEACON, J. W. Differential tropism to living and dead cereal root hairs by the biocontrol fungus <i>Idriella bolleyi</i>	217

Number 4, October 1992

ALAHAKOON, P. W., SREENIVASAPRASAD, S., BROWN, A. E. and MILLS, P. R. Selection of a genetic variant within <i>Colletotrichum gloeosporioides</i> isolates pathogenic on mango by passaging through wounded tomato fruits	227
PU, XIN-AN and GOODMAN, R. N. Induction of necrogenesis by <i>Agrobacterium tumefaciens</i> on grape explants	241
DICKMAN, M. B. and MITRA, A. <i>Arabidopsis thaliana</i> as a model for studying <i>Sclerotinia sclerotiorum</i> pathogenesis	255
SREENIVASAPRASAD, S., BROWN, AVERIL E. and MILLS, P. R. DNA sequence variation and interrelationships among <i>Colletotrichum</i> species causing strawberry anthracnose	265
LEE, S-S., KAWAKITA, K., TSUGE, T. and DOKE, N. Stimulation of phospholipase A ₂ activity in strawberry cells treated with AF-toxin I produced by <i>Alternaria alternata</i> strawberry pathotype	283
WHITEHEAD, DEBRA S., CODDINGTON, A. and LEWIS, B. G. Classification of races by DNA polymorphism analysis and vegetative compatibility grouping in <i>Fusarium oxysporum</i> f.sp. <i>pisi</i>	295

Number 5, November 1992

VIDHYASEKARAN, P., BORRAMEO, E. S. and MEW, T. W. <i>Helminthosporium oryzae</i> toxin suppresses phenol metabolism in rice plants and aids pathogen colonization	307
MANOCHA, M. S. and LONGCHENG SU. Immuno-cytochemical localization of host glycoproteins involved in the attachment of the mycoparasite <i>Piptocephalis virginiana</i>	317

BLOUNT, JACK W., DIXON, RICHARD A. and PAIVA, NANCY L. Stress responses in alfalfa (<i>Medicago sativa</i> L.) XVI. Antifungal activity of medicarpin and its biosynthetic precursors; implications for the genetic manipulation of stress metabolites	333
BENHAMOU, NICOLE. Ultrastructural detection of β -1,3-glucans in tobacco root tissues infected by <i>Phytophthora parasitica</i> var. <i>nicotianae</i> using a gold-complexed tobacco β -1,3-glucanase	351
CHÉRIF, MOHAMED, MENZIES, JAMES G., BENHAMOU, NICOLE and BELANGER, RICHARD R. Studies of silicon distribution in wounded and <i>Pythium ultimum</i> infected cucumber plants.. .. .	371

Number 6, December 1992

BALASS, M., COHEN, Y. and BAR-JOSEPH, M. Identification of a constitutive 45 kDa soluble protein associated with resistance to downy mildew in muskmelon (<i>Cucumis melo</i> L.), line PI 124111F	387
CARVER, T. L. W., ZEYEN, R. J., ROBBINS, M. P. and DEARNE, G. A. Effects of the PAL inhibitor, AOPP, on oat, barley and wheat cell responses to appropriate and inappropriate formae speciales of <i>Erysiphe graminis</i> DC	397
CHÉRIF, M., BENHAMOU, N., MENZIES, J. G. and BÉLANGER, R. R. Silicon induced resistance in cucumber plants against <i>Pythium ultimum</i>	411
MÖLLERS, C., ZITZLSPERGER, J. and WENZEL, G. The effects of a toxin preparation from <i>Phytophthora infestans</i> on potato protoplasts and microspores	427
WIJDEVELD, M. M. G., GOLDBACH, R. W., MEURS, C. and VAN LOON, L. C. Accumulation of viral 126 kDa protein and symptom expression in tobacco systemically infected with different strains of tobacco mosaic virus	437

Physiological and Molecular Plant Pathology

Index to Volume 41

- AF-toxin I produced by *Alternaria alternata* strawberry pathotype, stimulation of phospholipase A₂ activity in strawberry cells treated with, 283
- Agrobacterium tumefaciens*, induction of necrogenesis by, on grape explants, 241
- ALAHAKOON, P. W., SREENIVASAPRASAD, S., BROWN, A. E. and MILLS, P. R., Selection of a genetic variant within *Colletotrichum gloeosporioides* isolates pathogenic on mango by passaging through wounded tomato fruits, 227
- Alfalfa, stress responses; antifungal activity of medicarpin and its biosynthetic precursors; implications for genetic manipulation of stress metabolites, 333
- ALLAN, R. H., THORPE, C. J. and DEACON, J. W., Differential tropism to living and dead cereal root hairs by the biocontrol fungus *Idriella bolleyi*, 217
- Alternaria alternata* strawberry pathotype, stimulation of phospholipase A₂ activity in strawberry cells treated with AF-toxin I produced by, 283
- ANNIS, S. L., see XUE, B., 179
- Anthrachnose, strawberry, DNA sequence variation and interrelationships among *Colletotrichum* species causing, 265
- AOPP: effects on oat, barley and wheat cell responses to formae speciales of *Erysiphe graminis*, 397
- Arabidopsis thaliana* as a model for studying *Sclerotinia sclerotiorum* pathogenesis, 255
- Arachidonic acid: metabolism involved in eliciting activity in potato tuber, 127
- Arachidonic acid and its metabolites; evidence for release from sporangia of *Phytophthora infestans* during infection of potato, 61
- BAILEY, J. A., see PAIN, N. A., 111
- BALASS, M., COHEN, Y. and BAR-JOSEPH, M., Identification of a constitutive 45 kDa soluble protein associated with resistance to downy mildew in muskmelon (*Cucumis melo* L.), line PI 124111F, 387
- BAR-JOSEPH, M., see BALASS, M., 387
- Barley cell responses to formae speciales of *Erysiphe graminis*, effects of AOPP on, 397
- Barley, incompatibility conditioned by the *Mla* gene in powdery mildew of, 165
- BÉLANGER, R. R., see CHÉRIF, M., 371, 411
- BENHAMOU, N., Ultrastructural detection of β -1,3-glucans in tobacco root tissues infected by *Phytophthora parasitica* var. *nicotianae* using a gold-complexed tobacco β -1,3-glucanase, 351
- BENHAMOU, N. and THERIAULT, G., Treatment with chitosan enhances resistance of tomato plants to the crown and root rot pathogen *Fusarium oxysporum* f.sp. *radicis-lycopersici*, 33
- BENHAMOU, N., see CHÉRIF, M., 371, 411
- BLOUNT, J. W., DIXON, R. A. and PAIVA, N. L., Stress responses in alfalfa (*Medicago sativa* L.) XVI. Antifungal activity of medicarpin and its biosynthetic precursors; implications for the genetic manipulation of stress metabolites, 333
- BORROMEO, E. S., see VIDHYASEKARAN, P., 307
- BOSTOCK, R. M., see RICKER, K. E., 61
- BROWN, A. E., see ALAHAKOON, P. W., 227
- BROWN, A. E., see SREENIVASAPRASAD, S., 265
- BUSHNELL, W. R., see LEE-STADELMANN, O. Y., 165

- CARVER, T. L. W., ROBBINS, M. P., ZEYEN, R. J. and DEARNE, G. A., Effects of PAL-specific inhibition on suppression of activated defence and quantitative susceptibility of oats to *Erysiphe graminis*, 149
- CARVER, T. L. W., ZEYEN, R. J., ROBBINS, M. P. and DEARNE, G. A., Effects of the PAL inhibitor, AOPP, on oat barley and wheat cell responses to appropriate and inappropriate formae speciales of *Erysiphe graminis* DC, 397
- CASTLE, A. J., see RGHEI, N. A., 139
- CASTORIA, R., FANELLI, C., FABBRI, A. A. and PASSI, S., Metabolism of arachidonic acid involved in its eliciting activity in potato tuber, 127
- Cereal root hairs, living and dead, differential tropism to, by the biocontrol fungus *Idriella bolleyi*, 217
- CHÉRIF, M., BENHAMOU, N., MENZIES, J. G. and BÉLANGER, R. R., Silicon induced resistance in cucumber plants against *Pythium ultimum*, 411
- CHÉRIF, M., MENZIES, J. G., BENHAMOU, N. and BÉLANGER, R. R., Studies of silicon distribution in wounded and *Pythium ultimum* infected cucumber plants, 371
- Chitinase, resistance to diseases in the hybrid *Nicotiana glutinosa* × *Nicotiana debneyi* is associated with high levels of, 11
- Chitinases: subcellular localization in *Cladosporium fulvum* (syn. *Fulvia fulva*)-infected tomato leaves, 23
- Chitosan treatment enhances resistance of tomato plants to the crown and root rot pathogen *Fusarium oxysporum* f.sp. *radicis-lycopersici*, 33
- Cladosporium fulvum* (syn. *Fulvia fulva*)-infected tomato leaves, subcellular localization of plant chitinases and 1,3- β -glucanases in, 23
- CODDINGTON, A., see WHITEHEAD, D. S., 295
- COHEN, Y., see BALASS, M., 387
- Colletotrichum* species causing strawberry anthracnose, DNA sequence variation and interrelationships among, 265
- Colletotrichum gloeosporioides* isolates pathogenic on mango, selection of a genetic variant within, by passaging through wounded tomato fruits, 227
- Colletotrichum graminicola* elicitor: effects on the response of sorghum to *Periconia circinata* and its pathotoxin, 75
- Colletotrichum lindemuthianum*, monoclonal antibodies which show restricted binding to, 111
- Colletotrichum malvarum*, monoclonal antibodies which show restricted binding to, 111
- Colletotrichum orbiculare*, monoclonal antibodies which show restricted binding to, 111
- Colletotrichum trifolii*, monoclonal antibodies which show restricted binding to, 111
- Conidial exudate, cutinase is a component of; preparation of the infection court by *Erysiphe graminis* f.sp. *hordei*, 53
- COOPER, R. M., see FLOOD, J., 201
- CORDERO, M. J., RAVENTÓS, D. and SAN SEGUNDO, B., Induction of PR proteins in germinating maize seeds infected with the fungus *Fusarium moniliforme*, 189
- Cotton leaves, development of fluorescent, hypersensitively necrotic cells containing phytoalexins adjacent to colonies of *Xanthomonas campestris* pv. *malvacearum* in, 85
- Cotton leaves, a method for determining phytoalexin concentrations in fluorescent hypersensitively necrotic cells in, 101
- COVER, E. C., see ESSENBERG, M., 85, 101
- Cucumber plants, silicon induced resistance in, against *Pythium ultimum*, 411
- Cucumber plants, wounded and *Pythium ultimum* infected, silicon distribution in, 371
- Cucumis melo* L., line PI 124111F, identification of a 45 kDa soluble protein associated with resistance to downy mildew in, 387
- CURRAN, C. M., see LEE-STADELMANN, O. Y., 165
- Cutinase is a component of the conidial exudate; preparation of the infection court by *Erysiphe graminis* f.sp. *hordei*, 53
- DEACON, J. W., see ALLAN, R. H., 217
- DEARNE, G. A., see CARVER, T. L. W., 149, 397

- DE WIT, P. J. G. M., see WUBBEN, J. P., 23
- DICKMAN, M. B. and MITRA, A., *Arabidopsis thaliana* as a model for studying *Sclerotinia sclerotiorum* pathogenesis, 255
- DIXON, R. A., see BLOUNT, J. W., 333
- DNA polymorphism analysis and classification of races in *Fusarium oxysporum* f.sp. *pisi*, 295
- DNA polymorphisms and vegetative compatibility in *Fusarium oxysporum* f.sp. *elaeidis*; relationship to isolate virulence and origin, 201
- DNA sequence variation and interrelationships among *Colletotrichum* species causing strawberry anthracnose, 265
- DOKE, N., see LEE, S.-S., 283
- Downy mildew in muskmelon, line PI 124111F, identification of a 45 kDa soluble protein associated with resistance to, 387
- DUNKLE, L. D., see RANSOM, R. F., 75
- Elicitor from *Colletotrichum graminicola*: effects on the response of sorghum to *Periconia cinctata* and its pathotoxin, 75
- Erratum, 73
- Erysiphe graminis*, effects of AOPP on oat, barley and wheat cell responses to formae speciales of, 397
- Erysiphe graminis*, effects of PAL-specific inhibition on suppression of activated defence and quantitative susceptibility of oats to, 149
- Erysiphe graminis* f.sp. *hordei*, preparation of the infection court by; cutinase is a component of the conidial exudate, 53
- ESSENBERG, M., PIERCE, M. L., COVER, E. C., HAMILTON, B., RICHARDSON, P. E. and SCHOLES, V. E., A method for determining phytoalexin concentrations in fluorescent hypersensitively necrotic cells in cotton leaves, 101
- ESSENBERG, M., PIERCE, M. L., HAMILTON, B., COVER, E. C., SCHOLES, V. E. and RICHARDSON, P. E., Development of fluorescent, hypersensitively necrotic cells containing phytoalexins adjacent to colonies of *Xanthomonas campestris* pv. *malvacearum* in cotton leaves, 85
- FABBRI, A. A., see CASTORIA, R., 127
- FANELLI, C., see CASTORIA, R., 127
- FELIX, G., see GOY, P. A., 11
- Fimbriae: involvement in fungal host-mycoparasite interaction, 139
- FLOOD, J., WHITEHEAD, D. S. and COOPER, R. M., Vegetative compatibility and DNA polymorphisms in *Fusarium oxysporum* f.sp. *elaeidis* and their relationship to isolate virulence and origin, 201
- Fluorescent, hypersensitively necrotic cells containing phytoalexins adjacent to colonies of *Xanthomonas campestris* pv. *malvacearum* in cotton leaves, development of, 85
- Fluorescent hypersensitively necrotic cells in cotton leaves, a method for determining phytoalexin concentrations in, 101
- Fulvia fulva*, see *Cladosporium fulvum*, 23
- Fungal host-mycoparasite interaction, involvement of fimbriae in, 139
- Fusarium moniliforme*, induction of PR proteins in germinating maize seeds infected with, 189
- Fusarium oxysporum* f.sp. *elaeidis*, vegetative compatibility and DNA polymorphisms in, 201
- Fusarium oxysporum* f.sp. *pisi*, classification of races by DNA polymorphism analysis and vegetative compatibility grouping in, 295
- Fusarium oxysporum* f.sp. *radicis-lycopersici*, treatment with chitosan enhances resistance of tomato plants to, 33
- Gene *Mla*, incompatibility conditioned by, in powdery mildew of barley; change in permeability to non-electrolytes, 165
- Genetic manipulation of stress metabolites; antifungal activity of medicarpin; stress responses in alfalfa, 333

- Genetic variant selection within *Colletotrichum gloeosporioides* isolates pathogenic on mango by passaging through wounded tomato fruits, 227
- β -1,3-Glucanase, resistance to diseases in the hybrid *Nicotiana glutinosa* \times *Nicotiana debneyi* is associated with high levels of, 11
- 1,3- β -Glucanases: subcellular localization in *Cladosporium fulvum* (syn. *Fulvia fulva*)-infected tomato leaves, 23
- β -1,3-Glucans in tobacco root tissues infected by *Phytophthora parasitica* var. *nicotianae*; ultrastructural detection using a gold-complexed tobacco β -1,3-glucanase, 351
- Glycoproteins, host, involved in the attachment of the mycoparasite *Piptocephalis virginianam* immuno-cytochemical localization of, 317
- GOLDBACH, R. W., see WIJDEVELD, M. M. G., 437
- GOODMAN, R. N., see PU, XIN-AN, 241
- GOODWIN, P. H., see XUE, B., 179
- GORDON-WEEKS, R., see PIERPOINT, W. S., 1
- GOY, P. A., FELIX, G., MÉTRAUX, J. P. and MEINS, F. JR., Resistance to diseases in the hybrid *Nicotiana glutinosa* \times *Nicotiana debneyi* is associated with high constitutive levels of β -1,3-glucanase, chitinase, peroxidase and polyphenoloxidase, 11
- Grape explants, induction of necrogenesis by *Agrobacterium tumefaciens* on, 241
- GREEN, J. R., see PAIN, N. A., 111
- HAMILTON, B., see ESSENBERG, M., 85, 101
- Helminthosporium oryzae* toxin supresses phenol metabolism in rice plants and aids pathogen colonization, 307
- HIPSKIND, J., see RANSOM, R. F., 75
- Idriella bolleyi*, differential tropism to living and dead cereal root hairs by, 217
- Immuno-cytochemical localization of host glycoproteins involved in the attachment of the mycoparasite *Piptocephalis virginiana*, 317
- Intercellular fluids from *Nicotiana* species and from an interspecific *Nicotiana* hybrid, occurrence of protein PR-5 in, 1
- JACKSON, P. J., see PIERPOINT, W. S., 1
- JOOSTEN, M. H. A. J., see WUBBEN, J. P., 23
- KAWAKITA, K., see LEE, S-S., 283
- KUNOH, H., see PASCHOLATI, S. F., 53
- Leaves, cotton, development of fluorescent, hypersensitively necrotic cells containing phytoalexins adjacent to colonies of *Xanthomonas campestris* pv. *malvacearum* in, 85
- Leaves, cotton, a method for determining phytoalexin concentrations in fluorescent hypersensitively necrotic cells in, 101
- LEE, S-S., KAWAKITA, K., TSUGE, T. and DOKE, N., Stimulation of phospholipase A_2 activity in strawberry cells treated with AF-toxin I produced by *Alternaria alternata* strawberry pathotype, 283
- LEE-STADELMANN, O. Y., CURRAN, C. M. and BUSHNELL, W. R., Incompatibility conditioned by the *Mla* gene in powdery mildew of barley: change in permeability to non-electrolytes, 165
- LEITE, B., see RANSOM, R. F., 75
- Leptosphaeria maculans*, pathotype identification of, with PCR and oligonucleotide primers from ribosomal internal transcribed spacer sequences, 179
- LEWIS, B. G., see WHITEHEAD, D. S., 295
- LONGCHENG, SU., see MANOCHA, M. S., 317
- Maize seeds, germinating, infected with the fungus *Fusarium moniliforme*, induction of PR proteins in, 189

- Mango, selection of a genetic variant within *Colletotrichum gloeosporioides* isolates pathogenic on, by passing through wounded tomato fruits, 227
- MANOCHA, M. S. and LONGCHENG, SU., Immuno-cytochemical localization of host glycoproteins involved in the attachment of the mycoparasite *Piptocephalis virginiana*, 317
- MANOCHA, M. S., see RGHEI, N. A., 139
- Medicago sativa*, stress responses; antifungal activity of medicarpin and its biosynthetic precursors; implications for genetic manipulation of stress metabolites, 333
- Medicarpin antifungal activity; stress responses in alfalfa; implications for genetic manipulation of stress metabolites, 333
- MEINS, F. JR., see GOY, P. A., 11
- MENZIES, J. G., see CHÉRIF, M., 371, 411
- Metabolism of arachidonic acid involved in its eliciting activity in potato tuber, 127
- Metabolism, phenol, is suppressed by *Helminthosporium oryzae* toxin in rice plants, 307
- MÉTRAUX, J. P., see GOY, P. A., 11
- MEURS, C., see WIJDEVELD, M. M. G., 437
- MEW, T. W., see VIDHYASEKARAN, P., 307
- Microspores, potato, effects of a toxin preparation from *Phytophthora infestans* on, 427
- MILLS, P. R., see ALAHAKOON, P. W., 227
- MILLS, P. R., see SREENIVASAPRASAD, S., 265
- MITRA, A., see DICKMAN, M. B., 255
- Mla* gene, incompatibility conditioned by, in powdery mildew of barley; change in permeability to non-electrolytes, 165
- MÖLLERS, C., ZITZLSPERGER, J. and WENZEL, G., The effects of a toxin preparation from *Phytophthora infestans* on potato protoplasts and microspores, 427
- Monoclonal antibodies which show restricted binding to four *Colletotrichum* species: *C. lindemuthianum*, *C. malvarum*, *C. orbiculare* and *C. trifolii*, 111
- Muskmelon (*Cucumis melo* L.), line PI 124111F, identification of a 45 kDa protein associated with resistance to downy mildew in, 387
- Mycoparasite interaction with fungal host, involvement of fimbriae in, 139
- Mycoparasite *Piptocephalis virginiana*, immuno-cytochemical localization of host glycoproteins involved in the attachment of, 317
- Necrogenesis induction by *Agrobacterium tumefaciens* on grape explants, 241
- NICHOLSON, R. L., see PASCHOLATI, S. F., 53
- NICHOLSON, R. L., see RANSOM, R. F., 75
- Nicotiana* species and an interspecific *Nicotiana* hybrid, occurrence of protein PR-5 in intercellular fluids from, 1
- Nicotiana glutinosa* × *Nicotiana debneyi* hybrid, resistance to diseases in, 11
- Oat cell responses to formae speciales of *Erysiphe graminis*, effects of AOPP on, 397
- Oats: effects of PAL-specific inhibition on suppression of activated defence and quantitative susceptibility to *Erysiphe graminis*, 149
- O'CONNELL, R. J., see PAIN, N. A., 111
- Oligonucleotide primers from ribosomal internal transcribed spacer sequences, pathotype identification of *Leptosphaeria maculans* with PCR and, 179
- PAIN, N. A., O'CONNELL, R. J., BAILEY, J. A. and GREEN, J. R., Monoclonal antibodies which show restricted binding to four *Colletotrichum* species: *C. lindemuthianum*, *C. malvarum*, *C. orbiculare* and *C. trifolii*, 111
- PAIVA, N. L., see BLOUNT, J. W., 333
- PAL inhibitor, AOPP: effects on oat, barley and wheat cell responses to formae speciales of *Erysiphe graminis*, 397
- PAL-specific inhibition: effects on suppression of activated defence and quantitative susceptibility of oats to *Erysiphe graminis*, 149
- PASCHOLATI, S. F., YOSHIOKA, H., KUNOH, H. and NICHOLSON, R. L., Preparation of the

- infection court by *Erysiphe graminis* f.sp. *hordei*: cutinase is a component of the conidial exudate, 53
- PASSI, S., see CASTORIA, R., 127
- PCR and oligonucleotide primers from ribosomal internal transcribed spacer sequences, pathotype identification of *Leptosphaeria maculans* with, 179
- Periconia circinata*, effects of elicitor from *Colletotrichum graminicola* on the response of sorghum to, 75
- Peroxidase, resistance to diseases in the hybrid *Nicotiana glutinosa* × *Nicotiana debneyi* is associated with high levels of, 11
- Phenol metabolism in rice plants is suppressed by *Helminthosporium oryzae* toxin, 307
- Phospholipase A₂ activity stimulation in strawberry cells treated with AF-toxin I produced by *Alternaria alternata* strawberry pathotype, 283
- Phytoalexin concentrations in fluorescent hypersensitively necrotic cells in cotton leaves, a method for determination of, 101
- Phytoalexins, development of fluorescent, hypersensitively necrotic cells containing, adjacent to colonies of *Xanthomonas campestris* pv. *malvacearum* in cotton leaves, 85
- Phytophthora infestans*, evidence for release of the elicitor arachidonic acid and its metabolites from sporangia of, during infection of potato, 61
- Phytophthora infestans* toxin preparation; effects on potato protoplasts and microspores, 427
- Phytophthora parasitica* var. *nicotianae*, ultrastructural detection of β -1,3-glucans in tobacco root tissues infected by, 351
- PIERCE, M. L., see ESSENBERG, M., 85, 161
- PIERPPOINT, W. S., GORDON-WEEKS, R. and JACKSON, P. J., The occurrence of the thaumatin-like, pathogenesis-related protein, PR-5, in intercellular fluids from *Nicotiana* species and from an interspecific *Nicotiana* hybrid, 1
- Piptocephalis virginiana*, immuno-cytochemical localization of host glycoproteins involved in the attachment of, 317
- Polyphenoloxidase, resistance to diseases in the hybrid *Nicotiana glutinosa* × *Nicotiana debneyi* is associated with high levels of, 11
- Potato, evidence for release of elicitor arachidonic acid and its metabolites from sporangia of *Phytophthora infestans* during infection of, 61
- Potato protoplasts and microspores, effects of a toxin preparation from *Phytophthora infestans* on, 427
- Potato tuber, metabolism of arachidonic acid involved in its eliciting activity in, 127
- Powdery mildew of barley, incompatibility conditioned by the *Mla* gene in, 165
- PR proteins: induction in germinating maize seeds infected with the fungus *Fusarium moniliforme*, 189
- Protein, 126 kDa, viral, accumulation in tobacco systemically infected with different strains of tobacco mosaic virus, 437
- Protein PR-5: occurrence in intercellular fluids from *Nicotiana* species and from an interspecific *Nicotiana* hybrid, 1
- Protein, soluble, 45 kDa, associated with resistance to downy mildew in muskmelon, line PI 124111F, identification of, 387
- Protoplasts, potato, effects of a toxin preparation from *Phytophthora infestans* on, 427
- PU, XIN-AN and GOODMAN, R. N., Induction of necrogenesis by *Agrobacterium tumefaciens* on grape explants, 241
- Pythium ultimum* infected cucumber plants, silicon distribution in, 371
- Pythium ultimum*, silicon induced resistance in cucumber plants against, 411
- Race classification by DNA polymorphism analysis and vegetative compatibility grouping in *Fusarium oxysporum* f.sp. *pisi*, 295
- RANSOM, R. F., HIPSKIND, J., LEITE, B., NICHOLSON, R. L. and DUNKLE, L. D., Effects of elicitor from *Colletotrichum graminicola* on the response of sorghum to *Periconia circinata* and its pathotoxin, 75
- RAVENTÓS, D., see CORDERO, M. J., 189

- Resistance to diseases in the hybrid *Nicotiana glutinosa* × *Nicotiana debneyi*, 11
- Resistance to downy mildew in muskmelon, line PI 123111F, identification of a 45 kDa soluble protein associated with, 397
- Resistance, silicon induced, in cucumber plants against *Pythium ultimum*, 411
- Resistance of tomato plants to the crown and root rot pathogen *Fusarium oxysporum* f.sp. *radicis-lycopersici* is enhanced by treatment with chitosan, 33
- RGHEI, N. A., CASTLE, A. J. and MANOCHA, M. S., Involvement of fimbriae in fungal host-mycoparasite interaction, 139
- Rice plants, *Helminthosporium oryzae* toxin suppresses phenol metabolism in, and aids pathogen colonization, 307
- RICHARDSON, P. E., see ESSENBERG, M., 85, 101
- RICKER, K. E. and BOSTOCK, R. M., Evidence for release of the elicitor arachidonic acid and its metabolites from sporangia of *Phytophthora infestans* during infection of potato, 61
- ROBBINS, M. P., see CARVER, T. L. W., 149, 397
- Root hairs, cereal, living and dead, differential tropism to, by the biocontrol fungus *Idriella bolleyi*, 217
- Root rot pathogen *Fusarium oxysporum* f.sp. *radicis-lycopersici*, treatment with chitosan enhances resistance of tomato plants to, 33
- Root, tobacco, tissues infected by *Phytophthora parasitica* var. *nicotianae*, ultrastructural detection of β -1,3-glucans in, 351
- SAN SEGUNDO, B., see CORDERO, M. J., 189
- SCHOLES, V. E., see ESSENBERG, M., 85, 101
- Sclerotinia sclerotiorum* pathogenesis studied using *Arabidopsis thaliana* as a model, 255
- Seeds, maize, germinating, infected with the fungus *Fusarium moniliforme*, induction of PR proteins in, 189
- Silicon distribution in wounded and *Pythium ultimum* infected cucumber plants, 371
- Silicon induced resistance in cucumber plants against *Pythium ultimum*, 411
- Sorghum: response to *Periconia circinata* and its pathotoxin; effects of elicitor from *Colletotrichum graminicola*, 75
- Sporangia of *Phytophthora infestans*, evidence for release of the elicitor arachidonic acid and its metabolites from, during infection of potato, 61
- SREENIVASAPRASAD, S., BROWN, A. E. and MILLS, P. R., DNA sequence variation and interrelationships among *Colletotrichum* species causing strawberry anthracnose, 265
- SREENIVASAPRASAD, S., see ALAHAKOON, P. W., 227
- Strawberry anthracnose, DNA sequence variation and interrelationships among *Colletotrichum* species causing, 265
- Strawberry cells treated with AF-toxin I produced by *Alternaria alternata* strawberry pathotype, stimulation of phospholipase A₂ activity in, 283
- Stress responses in alfalfa (*Medicago sativa*); antifungal activity of medicarpin and its biosynthetic precursors; implications for the genetic manipulation of stress metabolites, 333
- Subcellular localization of plant chitinases and 1,3- β -glucanases in *Cladosporium fulvum* (syn. *Fulvia fulva*)-infected tomato leaves, 23
- Thaumatococin-like, pathogenesis-related protein, PR-5; occurrence in intercellular fluids from *Nicotiana* species and from an interspecific *Nicotiana* hybrid, 1
- THERIAULT, G., see BENHAMOU, N., 33
- THORPE, C. J., see ALLAN, R. H., 217
- Tobacco mosaic virus, accumulation of viral 126 kDa protein and symptom expression in tobacco systemically infected with different strains of, 437
- Tobacco root tissues infected by *Phytophthora parasitica* var. *nicotianae*, ultrastructural detection of β -1,3-glucans in, 351
- Tomato fruits, wounded, selection of a genetic variant within *Colletotrichum gloeosporioides* isolates pathogenic on mango by passing through, 227

- Tomato plants: treatment with chitosan enhances resistance to the crown and root rot pathogen, *Fusarium oxysporum* f.sp. *radicis-lycopersici*, 33
- Tropism, differential, to living and dead cereal root hairs by the biocontrol fungus *Idriella bolleyi*, 217
- TSUGE, T., see LEE, S.-S., 283
- Tuber, potato, metabolism of arachidonic acid involved in its eliciting activity in, 127
- Ultrastructural detection of β -1,3-glucans in tobacco root tissues infected by *Phytophthora parasitica* var. *nicotianae* using a gold-complexed tobacco β -1,3-glucanase, 351
- VAN KAN, J. A. L., see WUBBEN, J. P., 23
- VAN LOON, L. C., see WIJDEVELD, M. M. G., 437
- Vegetative compatibility and DNA polymorphisms in *Fusarium oxysporum* f.sp. *elaedis* and their relationship to isolate virulence and origin, 201
- Vegetative compatibility grouping and classification of races in *Fusarium oxysporum* f.sp. *pisi*, 295
- Vidhyasekaran, P., Borromeo, E. S. and MEW, T. W., *Helminthosporium oryzae* toxin suppresses phenol metabolism in rice plants and aids pathogen colonization, 307
- Viral 126 kDa protein accumulation and symptom expression in tobacco systemically infected with different strains of tobacco mosaic virus, 437
- WIJDEVELD, M. M. G., GOLDBACH, R. W., MEURS, C. and VAN LOON, L. C., Accumulation of viral 126 kDa protein and symptom expression in tobacco systemically infected with different strains of tobacco mosaic virus, 437
- WENZEL, G., see MÖLLERS, C., 427
- Wheat cell responses to formae speciales of *Erysiphe graminis*, effects of AOPP on, 397
- WHITEHEAD, D. S., CODDINGTON, A. and LEWIS, B. G., Classification of races by DNA polymorphism analysis and vegetative compatibility grouping in *Fusarium oxysporum* f.sp. *pisi*, 295
- WHITEHEAD, D. S., see FLOOD, J., 201
- WUBBEN, J. P., JOOSTEN, M. H. A. J., VAN KAN, J. A. L. and DE WIT, P. J. G. M., Subcellular localization of plant chitinases and 1,3- β -glucanases in *Cladosporium fulvum* (syn. *Fulvia fulva*)-infected tomato leaves, 23
- Xanthomonas campestris* pv. *malvacearum* colonies in cotton leaves, development of fluorescent, hypersensitively necrotic cells containing phytoalexins adjacent to, 85
- XUE, B., GOODWIN, P. H. and ANNIS, S. L., Pathotype identification of *Leptosphaeria maculans* with PCR and oligonucleotide primers from ribosomal internal transcribed spacer sequences, 179
- YOSHIOKA, H., see PASCHOLATI, S. F., 53
- ZEYEN, R. J., see CARVER, T. L. W., 149, 397
- ZITZLSPERGER, J., see MÖLLERS, C., 427